arXiv:0706.1077v1 [math.AP] 7 Jun 2007

REGULARITY OF DIRICHLET NEARLY MINIMIZING MULTIPLE-VALUED FUNCTIONS

JORDAN GOBLET AND WEI ZHU

Abstract. In this paper, we extend the related notions of Dirichlet quasiminimizer, $\omega-$minimizer and almost minimizer to the framework of multiple-valued functions in the sense of Almgren and prove Hölder regularity results. We also give examples of those minimizers with various branch sets.

Introduction

In [2] F. J. Almgren, Jr. has developed a theory of functions defined on a bounded open subset Ω of $\mathbb{R}^m$ with smooth boundary and taking values in the space $\mathbf{Q}_Q(\mathbb{R}^n)$ of unordered Q tuples of points of $\mathbb{R}^n$ (see [1] for a summary). In terms of Dirichlet's integral naturally defined for appropriate functions $\Omega \to \mathbf{Q}_Q(\mathbb{R}^n)$ he obtained the following existence and regularity results:

Theorem. *For each appropriate function $v : \partial\Omega \to \mathbf{Q}_Q(\mathbb{R}^n)$ there exists a function $u : \Omega \to \mathbf{Q}_Q(\mathbb{R}^n)$ having boundary values v and of least Dirichlet integral among such functions. Furthermore, each such minimizer u is Hölder continuous.*

Almgren introduced the machinery of Dirichlet minimizing multiple-valued functions in order to study the partial regularity of mass-minimizing integral currents. He proved that any m-dimensional mass-minimizing integral current is regular except on a set of Hausdorff dimension at most $m-2$. In the present paper, we will define and study the regularity of Dirichlet quasiminimizer as well as $\omega-$minimizer and almost minimizer in the setting of multiple-valued functions.

The notion of quasiminimizer, introduced by M. Giaquinta and E. Giusti [7], embodies the notion of minimizer of variational integrals

$$J(u,\Omega) = \int_\Omega f(x,u,\mathrm{D}u)\,dx \tag{1}$$

where Ω is an open subset of $\mathbb{R}^m$, $u : \Omega \to \mathbb{R}^n$, $\mathrm{D}u = \{\mathrm{D}_\alpha u^i\}$, $\alpha = 1,\dots,m$, $i = 1,\dots,n$ and $f(x,u,p) : \Omega \times \mathbb{R}^n \times \mathbb{R}^{mn} \to \mathbb{R}$, but it is substantially more general. By definition, a quasiminimizer u minimizes the functional J only up to a multiplicative constant, that is, there exists $K \geq 1$ such that

$$J(u,V) \leq K\ J(u+\phi,V)$$

for each subdomain V that is compactly contained in Ω and for each $\phi \in C_0^\infty(V)$. In the scalar-valued case and under suitable conditions on f, Hölder continuity can be

---

Date: February 1, 2008.

Key words and phrases. Multiple-valued function, quasiminimizer, ω-minimizer, almost minimizer.

derived directly from the quasiminimizing property. For more details of the statement above, as well as for more information on the properties of quasiminimizers, we refer to [6], [7] and [8].

An alternative notion called ω-minimizer is considered in a paper by G. Anzellotti [3]. A function $u \in W^{1,2}(\Omega, \mathbb{R}^n)$ is a ω-minimizer, where $\omega : (0, R_0) \to \mathbb{R}$ is a nonnegative function, for a functional J of the type (1) if one has

$$J(u, U^m(x,r)) \leq (1 + \omega(r)) \; J(u + \phi, U^m(x,r))$$

for any ball $U^m(x,r) \subset \Omega$ with $r < R_0$ and for all functions $\phi \in W^{1,2}_0(U^m(x,r), \mathbb{R}^n)$. If the functional J is the Dirichlet integral and $\lim_{r \to 0} \omega(r) = 0$ then Anzellotti proves that u is locally Hölder continuous. In addition the first derivatives of u are locally Hölder continuous, with exponent γ, in Ω provided $0 \leq \omega(r) \leq cr^{2\gamma}$.

Motivated by the question of boundary regularity of Dirichlet minimizing multiple-valued functions (more precisely, subtracting a suitable extension of boundary value from the original minimizer gives an almost minimizer), we also consider (c, α)-almost minimizer in the following sense:

$$J(u, U^m(x,r)) \leq J(v, U^m(x,r)) + c \, r^{m-2+\alpha}$$

for any ball $U^m(x,r) \subset \Omega$ and for all functions $v \in W^{1,2}(U^m(x,r), \mathbb{R}^n)$ such that $u = v$ on $\partial U^m(x,r)$. Readers are also referred to [10] for the discussion of almost minimizer in the study of ferromagnetism in $\mathbb{R}^3$.

This work is organized as follows. In Section 2 we define the notion of Dirichlet quasiminimizing multiple-valued functions and Hölder regularity results are derived from two Almgren's estimates. We also construct a one-dimensional quasiminimizer with a fractal branch set. In Section 3 we prove a similar result to Anzellotti's Theorem for ω-minimizing multiple-valued functions and we give an example of $\omega-$minimizer with a unique branch point. In Section 4 we prove a Hölder regularity theorem for Dirichlet almost minimizing multiple-valued functions and construct a one-dimensional almost minimizer with a fractal branch set.

1. Preliminaries

In our terminology and notations we shall follow scrupulously [2] and [5]. Throughout the whole text m, n and Q will be positive integers. We denote by $U^m(x,r)$ and $B^m(x,r)$ the open and closed balls in $\mathbb{R}^m$ with center x and radius r.

1.1. The metric space $\mathbf{Q}_Q(\mathbb{R}^n)$.

The space of unordered Q tuples of points in $\mathbb{R}^n$ is

$$\mathbf{Q}_Q(\mathbb{R}^n) = \left\{ \sum_{i=1}^{Q} [[u_i]] : u_1, \ldots, u_Q \in \mathbb{R}^n \right\},$$

where $[[u_i]]$ is the Dirac measure at u_i. This space is equipped with the metric $\mathcal{G}$;

$$\mathcal{G}\left(\sum_{i=1}^{Q} [[u_i]], \sum_{i=1}^{Q} [[v_i]] \right) = \min_{\sigma} \left(\sum_{i=1}^{Q} |u_i - v_{\sigma(i)}|^2 \right)^{1/2}$$

where σ runs through all the permutations of $\{1, \ldots, Q\}$. A multiple-valued function in the sense of Almgren is a $\mathbf{Q}_Q(\mathbb{R}^n)$-valued function.

There are a bi-Lipschitz homeomorphism $\xi : \mathbf{Q}_Q(\mathbb{R}^n) \to Q^* \subset \mathbb{R}^{PQ}$, where P is a positive integer, and a Lipschitz retraction $\rho : \mathbb{R}^{PQ} \to Q^*$ with $\rho|_{Q^*} = \mathrm{Id}_{Q^*}$ (see Theorems 1.2(3) and 1.3(1) in [2]).

1.2. **Affine approximations.** The set of all affine maps $\mathbb{R}^m \to \mathbb{R}^n$ is denoted by $A(m,n)$. If $v = (v_1, \ldots, v_n) \in A(m,n)$, we put

$$|v| = \left(\sum_{i=1}^{n} \sum_{j=1}^{m} \left(\frac{\partial v_i}{\partial x_j} \right)^2 \right)^{1/2} \in \mathbb{R}.$$

A function $v : \mathbb{R}^m \to \mathbf{Q}_Q(\mathbb{R}^n)$ is called affine if there are $v_1, \ldots, v_Q \in A(m,n)$ such that $v = \sum_{i=1}^{Q} [[v_i]]$. Then we set $|v| = \left(\sum_{i=1}^{Q} |v_i|^2 \right)^{1/2}$.

If $\Omega \subset \mathbb{R}^m$ is open and $a \in \Omega$, a function $u : \Omega \to \mathbf{Q}_Q(\mathbb{R}^n)$ is said to be approximately affinely approximable at a if there is an affine function $v : \mathbb{R}^m \to \mathbf{Q}_Q(\mathbb{R}^n)$ such that

$$\operatorname{ap} \lim_{x \to a} \frac{\mathcal{G}(u(x), v(x))}{|x - a|} = 0.$$

Such a function v is uniquely determined and denoted by $\operatorname{ap}Au(a)$. This is the case if $\xi \circ u$ is approximately differentiable at a, and then

$$(\operatorname{Lip}(\xi))^{-1} |\operatorname{ap}D(\xi \circ u)(a)| \le |\operatorname{ap}Au(a)| \le |\operatorname{ap}D(\xi \circ u)(a)|$$

(see Theorem 1.4(3) in [2]).

1.3. **The Sobolev space $\mathcal{Y}_2(U, \mathbf{Q}_Q(\mathbb{R}^n))$.** Suppose U is an open ball in $\mathbb{R}^m$ or the entire space $\mathbb{R}^m$.

(1) We denote by $W^{1,2}(U, \mathbb{R}^n)$ the Sobolev space of $\mathbb{R}^n$-valued functions on U which together with their first order distributional partial derivatives are $\mathcal{L}^m$ square summable over U.

(2) A function $u \in W^{1,2}(U, \mathbb{R}^n)$ is said to be strictly defined if $u(x) = y$ whenever $x \in U, y \in \mathbb{R}^n$ and

$$\lim_{r \to 0} r^{-m} \int_{B^m(x,r)} |u(z) - y| \, d\mathcal{L}^m z = 0.$$

Any $v \in W^{1,2}(U, \mathbb{R}^n)$ agrees $\mathcal{L}^m$ almost everywhere on U with a strictly defined $u \in W^{1,2}(U, \mathbb{R}^n)$. If $u \in W^{1,2}(U, \mathbb{R}^n)$ is strictly defined, then

$$\lim_{r \to 0} r^{-m} \int_{B^m(x,r)} |u(z) - u(x)| \, d\mathcal{L}^m z = 0$$

for $\mathcal{H}^{m-1}$ almost all $x \in U$ (see Section 4.8 in [4]).

(3) The space $\mathcal{Y}_2(U, \mathbf{Q}_Q(\mathbb{R}^n))$ consists of those functions $u : U \to \mathbf{Q}_Q(\mathbb{R}^n)$ for which $\xi \circ u \in W^{1,2}(U, \mathbb{R}^{PQ})$. We say that u is strictly defined if $\xi \circ u$ is strictly defined. If $u, v \in \mathcal{Y}_2(U, \mathbf{Q}_Q(\mathbb{R}^n))$, we write $u = v$ if $u(x) = v(x)$ for $\mathcal{L}^m$ almost all $x \in U$.

(4) Suppose $u \in \mathcal{Y}_2(U, \mathbf{Q}_Q(\mathbb{R}^n))$. Then u is approximately affinely approximable $\mathcal{L}^m$ almost everywhere on U since $\xi \circ u$ is (see Section 6.1.3 in [4]). The Dirichlet integral of u over U is defined by

$$\operatorname{Dir}(u; U) = \int_U |\operatorname{ap}Au(x)|^2 \, d\mathcal{L}^m x.$$

1.4. **The Sobolev space $\partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$.** Suppose $U \subset \mathbb{R}^m$ is an open ball.

(1) The space $\partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ will be the set of those functions $v : \partial U \to \mathbf{Q}_Q(\mathbb{R}^n)$ for which there is a strictly defined $u \in \mathcal{Y}_2\left(\mathbb{R}^m, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ such that $u(x) = v(x)$ for $\mathcal{H}^{m-1}$ almost all $x \in \partial U$ (cf. Theorem A.1.2(7) and Section 2.1(2) in [2]). If $v, w \in \partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$, we write $v = w$ if $v(x) = w(x)$ for $\mathcal{H}^{m-1}$ almost all $x \in \partial U$.

(2) If $v \in \partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ we say that u has boundary values v if there is $w \in \mathcal{Y}_2\left(\mathbb{R}^m, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ which is strictly defined such that $w|_U = u|_U$ and $w|_{\partial U} = v$. We then write $u|_{\partial U} = v$. If u is strictly defined, v agrees with u $\mathcal{H}^{m-1}$ almost everywhere on ∂U.

(3) One says that $u : U \to \mathbf{Q}_Q(\mathbb{R}^n)$ is Dirichlet minimizing if and only if $u \in \mathcal{Y}_2\left(U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ and, assuming u has boundary values $v \in \partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$, one has

$$\mathrm{Dir}\,(u; U) = \inf\left\{\mathrm{Dir}\,(w; U) : w \in \mathcal{Y}_2\left(U, \mathbf{Q}_Q(\mathbb{R}^n)\right) \text{ has boundary values } v\right\}.$$

If $v \in \partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ then there exists a Dirichlet minimizing $u \in \mathcal{Y}_2\left(U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$ that has boundary values v (see Theorem 2.2(2) in [2]).

(4) For each $v \in \partial\mathcal{Y}_2(\partial U, \mathbf{Q}_Q(\mathbb{R}^n))$, the Dirichlet integral of v over ∂U is defined by

$$\mathrm{dir}(v; \partial U) = \int_{\partial U} |\mathrm{ap}Av(x)|^2 \, d\mathcal{H}^{m-1}x$$

where the affine approximation $\mathrm{ap}Av$ is with respect to ∂U.

2. Dirichlet quasiminimizer

Throughout this section K will be a real number larger than 1.

Definition 1. *A strictly defined function $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ is a Dirichlet K-quasiminimizer if for every ball $U \subset U^m(0,1)$,*

$$Dir(u; U) \leq K \; Dir(v; U)$$

where v is a Dirichlet minimizing multiple-valued function having boundary values $u|_{\partial U} \in \partial\mathcal{Y}_2\left(\partial U, \mathbf{Q}_Q(\mathbb{R}^n)\right)$.

2.1. **Regularity.** For the convenience of the reader, we shall recall here a few known results that we are going to need later. We begin with a "Dirichlet growth" theorem guaranteeing Hölder continuity:

Theorem 3.5.2 in [15]. *Suppose $u \in W^{1,p}(B^m(x_0, R))$, $0 < \sigma < 1$ and*

$$\int_{B(z,r)} |\nabla u(x)|^p \, dx \leq C \, (r/\delta)^{m-p+p\sigma}, \quad 0 \leq r \leq \delta = R - |z - x_0|$$

for every $z \in B(x_0, R)$. Then

$$u \in C^{0,\sigma}(B^m(x_0, r))$$

for each $r < R$.

We recall Almgren's estimates for 2-dimensional and higher dimensional domains:

Section 2.7(2) in [2]. *Suppose $v \in \partial\mathcal{Y}_2\left(\partial U^2(x,r), \mathbf{Q}_Q(\mathbb{R}^n)\right)$. Then there exists a multiple-valued function $u \in \mathcal{Y}_2\left(U^2(x,r), \mathbf{Q}_Q(\mathbb{R}^n)\right)$ such that*

(1) *u has boundary values v,*

(2) $Dir\big(u;U^2(x,r)\big) \le Qr\ dir\big(v;\partial U^2(x,r)\big)$.

Theorem 2.12 in [2]**.** *Let $m \ge 3$ be an integer. If $f \in \mathcal{Y}_2(U^m(x,r),\mathbf{Q}_Q(\mathbb{R}^n))$ is strictly defined and Dirichlet minimizing then*

$$Dir(f;U^m(x,r)) \le \left(\frac{1-\varepsilon_Q}{m-2}\right) r\ dir(f;\partial U^m(x,r))$$

where $0<\varepsilon_Q<1$ is a constant.

Theorem 2. *Suppose that $u \in \mathcal{Y}_2\left(U^2(0,1),\mathbf{Q}_Q(\mathbb{R}^n)\right)$ is a strictly defined Dirichlet K-quasiminimizer such that $Dir\left(u;U^2(0,1)\right) > 0$. Then the following assertions are checked:*

1. *Suppose $z \in U^2(0,1)$. Then for $\mathcal{L}^1$ almost all $0<r<1-|z|$,*
$$Dir(u;U^2(z,r)) \le KQr\,\frac{d}{ds}\,Dir(u;U^2(z,s))|_{s=r}.$$
2. *For each $z \in U^2(0,1)$, $0<r<1-|z|$, and $0<s\le 1$,*
$$Dir(u;U^2(z,sr)) \le s^{\frac{1}{KQ}}\ Dir(u;U^2(z,r)).$$
3. *$f|_{B^2(0,\delta)}$ is Hölder continuous with exponent $\frac{1}{2KQ}$ for all $0<\delta<1$.*

Proof. Let $v \in \mathcal{Y}_2(U^2(z,r),\mathbf{Q}_Q(\mathbb{R}^n))$ be a Dirichlet minimizing multiple-valued function with boundary values $u|_{\partial U^2(z,r)}$. It follows from Section 2.7(2) in [2] that

$$\mathrm{Dir}(u;U^2(z,r)) \le K\ \mathrm{Dir}(v;U^2(z,r)) \le KQr\ \mathrm{dir}(u;\partial U^2(z,r)).$$

Using the fact that the function $\phi(r)$ $=$Dir$(u;U^2(z,r))$ is absolutely continuous with

$$\phi'(r) = \int_{\partial U^2(z,r)} |\mathrm{ap}Au(x)|^2\ d\mathcal{H}^1 x \ge \mathrm{dir}(u;\partial U^2(z,r))$$

for $\mathcal{L}^1$ almost all $0<r<1-|z|$, we obtain the first assertion. One derives (2) from (1) by integration. In order to verify (3), one notes that conclusion (2) implies for each $z \in U^2(0,1)$, $0<r<1-|z|$, and $0<s\le 1$,

$$\begin{aligned}\mathrm{Dir}(\xi\circ u;U^2(z,sr)) &\le \mathrm{Lip}(\xi)^2\ \mathrm{Dir}(u;U^2(z,sr))\\ &\le \mathrm{Lip}(\xi)^2\ s^{\frac{1}{KQ}}\ \mathrm{Dir}(u;U^2(z,r))\\ &\le \mathrm{Lip}(\xi)^2\ s^{\frac{1}{KQ}}\ \mathrm{Dir}(\xi\circ u;U^2(z,r))\\ &\le \mathrm{Lip}(\xi)^2\ \mathrm{Dir}(\xi\circ u;U^2(0,1))\ s^{\frac{1}{KQ}}.\end{aligned}$$

Let us fix $0<\delta<1$. Theorem 3.5.2 in [15] applied to the function $\xi\circ u$ implies that

$$\xi\circ u \in C^{0,\frac{1}{2KQ}}(B^2(0,\delta))$$

and the third assertion is checked since ξ is bi-Lipschitz. □

The regularity result is weaker for higher dimensional domains since we need an extra assumption (2) on the size of K. This assumption on K is necessary since M. Giaquinta presents in [6] a single-valued quasiminimizer for the Dirichlet integral which is singular on a dense set.

Theorem 3. *Let $m \geq 3$ be an integer. Suppose that $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ is a strictly defined Dirichlet K-quasiminimizer such that Dir$(u; U^m(0,1)) > 0$. If*

$$1 \leq K < \frac{1}{(1-\varepsilon_Q)} \tag{2}$$

then there exists a constant $0 < \sigma = \sigma(K, \varepsilon_Q, m) < 1$ such that the following assertions are checked:

1. *Suppose $z \in U^m(0,1)$. Then for $\mathcal{L}^1$ almost all $0 < r < 1 - |z|$,*
$$Dir(u; U^m(z,r)) \leq \frac{1}{(m-2+2\sigma)}\, r\, \frac{d}{ds}\, Dir(u; U^m(z,s))|_{s=r}.$$
2. *For each $z \in U^m(0,1)$, $0 < r < 1 - |z|$, and $0 < s \leq 1$,*
$$Dir(u; U^m(z,sr)) \leq s^{m-2+2\sigma}\, Dir(u; U^m(z,r)).$$
3. *$f|_{B^m(0,\delta)}$ is Hölder continuous with exponent σ for all $0 < \delta < 1$.*

Proof. We fix $0 < \sigma < 1$ requiring that

$$\frac{m-2}{K(1-\varepsilon_Q)} > m - 2 + 2\sigma.$$

Let $v \in \mathcal{Y}_2(U^m(z,r), \mathbf{Q}_Q(\mathbb{R}^n))$ be a Dirichlet minimizing multiple-valued function with boundary values $u|_{\partial U^m(z,r)}$. It follows from Theorem 2.12 in [2] that

$$\begin{aligned}
\mathrm{Dir}(u; U^m(z,r)) &\leq K\, \mathrm{Dir}(v; U^m(z,r)) \\
&\leq K \left(\frac{1-\varepsilon_Q}{m-2}\right) r\, \mathrm{dir}(v; \partial U^m(z,r)) \\
&\leq \frac{1}{(m-2+2\sigma)}\, r\, \mathrm{dir}(u; \partial U^m(z,r)).
\end{aligned}$$

Using the fact that the function $\phi(r) = \mathrm{Dir}(u; U^m(z,r))$ is absolutely continuous with

$$\phi'(r) = \int_{\partial U^m(z,r)} |\mathrm{ap}Au(x)|^2\, d\mathcal{H}^{m-1}x \geq \mathrm{dir}(u; \partial U^m(z,r))$$

for $\mathcal{L}^1$ almost all $0 < r < 1 - |z|$, we obtain the first claim. One derives (2) from (1) by integration. In order to conclude, one notes that the second claim implies for each $z \in U^m(0,1)$, $0 < r < 1 - |z|$, and $0 < s \leq 1$,

$$\begin{aligned}
\mathrm{Dir}(\xi \circ u; U^m(z,sr)) &\leq \mathrm{Lip}(\xi)^2\, \mathrm{Dir}(u; U^m(z,sr)) \\
&\leq \mathrm{Lip}(\xi)^2\, s^{m-2+2\sigma}\, \mathrm{Dir}(u; U^m(z,r)) \\
&\leq \mathrm{Lip}(\xi)^2\, s^{m-2+2\sigma}\, \mathrm{Dir}(\xi \circ u; U^m(z,r)) \\
&\leq \mathrm{Lip}(\xi)^2\, \mathrm{Dir}(\xi \circ u; U^m(0,1))\, s^{m-2+2\sigma}.
\end{aligned}$$

Let us fix $0 < \delta < 1$. Theorem 3.5.2 in [15] applied to the function $\xi \circ u$ implies that

$$\xi \circ u \in C^{0,\sigma}(B^m(0,\delta))$$

and the third assertion is checked since ξ is bi-Lipschitz. □

2.2. **Branch set of quasiminimizer.**

Definition 4. *Assume the multiple-valued function* $u : U^m(0,1) \to \mathbf{Q}_Q(\mathbb{R}^n)$ *is continuous. Define the function* $\sigma : U^m(0,1) \to \{1,\dots,Q\}$ *as*

$$\sigma(x) = \mathit{card}\ (\mathit{spt}\ (u(x)))$$

for all $x \in U^m(0,1)$. *Then, the closed set*

$$\Sigma = \{x \in U^m(0,1) \mid \sigma \text{ is discontinuous at } x\}$$

is called the branch set of the continuous multiple-valued function u.

Almgren proved that the Hausdorff dimension of the branch set of a strictly defined and Dirichlet minimizing $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ does not exceed $m-2$ (see Theorem 2.14 in [2]). Here we will construct a one-dimensional Dirichlet quasiminimizer with a fractal branch set. This construction can be adapted in order to obtain a one-dimensional Dirichlet quasiminimizer with a branch set of positive 1-dimensional Lebesgue measure.

The following result characterizes Dirichlet minimizing multiple-valued functions in codimension one.

Theorem 2.16 in [2]**.** *Suppose* $v \in \partial\mathcal{Y}_2(\partial U^m(0,1), \mathbf{Q}_Q(\mathbb{R}))$ *and* $v_1, \dots, v_Q : \partial U^m(0,1) \to \mathbb{R}$ *are defined by requiring for each* $x \in \partial U^m(0,1)$ *that* $v(x) = \sum_{i=1}^Q [[v_i(x)]]$ *and* $v_1(x) \le v_2(x) \le \dots \le v_Q(x)$. *Then* $v_i \in \partial\mathcal{Y}_2(\partial U^m(0,1), \mathbb{R})$ *for each* $i = 1, \dots, Q$ *and* $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}))$ *is strictly defined and Dirichlet minimizing with boundary values* v *if and only if* $u = \sum_{i=1}^Q [[u_i]]$ *corresponding to* $u_1, u_2, \dots, u_Q \in \mathcal{Y}_2(U^m(0,1), \mathbb{R})$ *which are the unique harmonic functions having boundary values* $v_1, v_2, \dots, v_Q$ *respectively.*

This last theorem implies the following characterization of Dirichlet quasiminimizers in dimension and codimension one.

Theorem 5. *A multiple-valued function* $u \in \mathcal{Y}_2((-1,1), \mathbf{Q}_Q(\mathbb{R}))$ *is a Dirichlet K-quasiminimizer if and only if for every interval* $(a,b) \subset (-1,1)$,

$$\mathit{Dir}(u, (a,b)) \le K\, \frac{\mathcal{G}^2(u(b), u(a))}{b-a}.$$

Proof. Let $(a,b) \subset (-1,1)$. We define the functions $v_1, \dots, v_Q : \partial(a,b) \to \mathbb{R}$ by requiring for each $x \in \partial(a,b)$ that $u(x) = \sum_{i=1}^Q [[v_i(x)]]$ and $v_1(x) \le v_2(x) \le \dots \le v_Q(x)$. By Theorem 2.16 in [2], the Dirichlet minimizing multiple-valued function coinciding with u on $\partial(a,b)$ is given by

$$v(\cdot) = \sum_{i=1}^{Q} \left[\left[\frac{v_i(b) - v_i(a)}{b-a}(\cdot - a) + v_i(a) \right]\right]$$

and

$$\begin{aligned}\mathrm{Dir}(v; [a,b]) = \int_a^b |\mathrm{ap}Av(x)|^2\, dx = \frac{(b-a)}{(b-a)^2} \sum_{i=1}^{Q} |v_i(b) - v_i(a)|^2 \\ = \frac{\mathcal{G}^2(u(b), u(a))}{b-a}.\end{aligned}$$

□

We will now produce a Dirichlet quasiminimizer with a fractal branch set. We begin with some definitions:

1. A *diamond* above an interval $I = [a, b] \subset \mathbb{R}$ is a multiple-valued function $u : [a, b] \to \mathbf{Q}_2(\mathbb{R})$ whose the graph is a parallelogram with the following vertices $[a, h], [(a+b)/2, h], [(a+b)/2, h+(b-a)/2], [b, h+(b-a)/2]$ where $h \in \mathbb{R}$.
2. A *pluri-diamond* is a multiple-valued function $u : [0, 1] \to \mathbf{Q}_2(\mathbb{R})$ which admits a partition of $[0, 1]$ in intervals $\{I_j\}$ such that
   - u is a diamond above I_j or a map of the form $x \to 2[[x+p_j]]$ above I_j where $p_j \in \mathbb{R}$,
   - u is continuous on $[0, 1]$.

Figure 1 gives examples of pluri-diamonds. It is clear that a pluri-diamond u is Lipschitz with $\mathrm{Lip}(u) \le \sqrt{2}$. We record two lemmas for later use:

Lemma 6. *A pluri-diamond u is a Dirichlet 4-quasiminimizer.*

Proof. Let $u_{\min}, u_{\max} : [0, 1] \to \mathbb{R}$ be such that $u = [[u_{\min}]] + [[u_{\max}]]$ and $u_{\min}(x) \le u_{\max}(x)$ for all $x \in [0, 1]$. For every interval $[a, b] \subset [0, 1]$, we have

$$\begin{aligned}\mathcal{G}^2(u(b), u(a)) &\ge \frac{1}{2}\left(|u_{\min}(b) - u_{\min}(a)| + |u_{\max}(b) - u_{\max}(a)|\right)^2 \\ &= \frac{1}{2}\left(\int_a^b u'_{\min}(x) + u'_{\max}(x)\, dx\right)^2 \ge \frac{(b-a)^2}{2}\end{aligned}$$

and

$$\begin{aligned}\frac{(b-a)\int_a^b |\mathrm{ap}Au(x)|^2\, dx}{\mathcal{G}^2(u(b), u(a))} &= \frac{(b-a)\int_a^b u'^2_{\min}(x) + u'^2_{\max}(x)\, dx}{\mathcal{G}^2(u(b), u(a))} \\ &\le \frac{2\,(b-a)^2}{\mathcal{G}^2(u(b), u(a))} \le 4\end{aligned}$$

hence u is a Dirichlet 4-quasiminimizer by Theorem 5. □

Lemma 23.7 in [17]. *Let Ω be an open subset of $\mathbb{R}^m$, $1 < p < \infty$. If $\{u_i\}$ is a bounded sequence in $W^{1,p}(\Omega)$ such that $u_i \rightharpoonup u$ in $L^p(\Omega)$ then $u \in W^{1,p}(\Omega)$ and*

$$\|\nabla u\|_p \le \liminf_{i\to\infty} \|\nabla u_i\|_p.$$

We will now create a particular sequence of pluri-diamonds $\{u^i\}$. We define $u^1 : \mathbb{R} \to \mathbf{Q}_2(\mathbb{R})$ by requiring that

- u^1 is a pluri-diamond,
- $u^1(1/3) = 0$,
- u^1 is a diamond only above $[1/3, 2/3]$.

We define $u^2 : \mathbb{R} \to \mathbf{Q}_2(\mathbb{R})$ by requiring that

- u^2 is a pluri-diamond,
- $u^2(1/3) = 0$,
- u^2 is a diamond only above the intervals $[1/3, 2/3]$, $[1/9, 2/9]$ and $[7/9, 8/9]$.

The reader will easily imagine how the remaining terms of the sequence are defined. For each $i \in \mathbb{N}$, we define $u^i_{\min}, u^i_{\max} : [0, 1] \to \mathbb{R}$ by requiring that

$$u^i(x) = [[u^i_{\min}(x)]] + [[u^i_{\max}(x)]] \quad \text{and} \quad u^i_{\min}(x) \le u^i_{\max}(x)$$



FIGURE 1. The pluri-diamonds u^1, u^2 and u^3.

for all $x \in [0,1]$. One can checks that

$$\mathrm{Lip}(u^i_{\min}) = \mathrm{Lip}(u^i_{\max}) = 1$$

for each $i \in \mathbb{N}$. By Ascoli's Theorem, we can extract two subsequences still denoted $\{u^i_{\min}\}$ and $\{u^i_{\max}\}$ and there exists two 1-Lipschitz functions $u_{\min}, u_{\max} : [0,1] \to \mathbb{R}$ such that $\{u^i_{\min}\}$ and $\{u^i_{\max}\}$ converge uniformly to $u_{\min}$ and $u_{\max}$. We define the Lipschitz multiple-valued function $u : [0,1] \to \mathbf{Q}_2(\mathbb{R})$ by

$$u = [[u_{\min}]] + [[u_{\max}]]. \tag{3}$$

Proposition 7. *The multiple-valued function u defined by (3) is a Dirichlet 4-quasiminimizer and the branch set of u is the Cantor ternary set.*

Proof. We estimate for any interval $[a,b] \subset [0,1]$,

$$\begin{aligned}
\int_a^b |\mathrm{ap}Au(x)|^2 \, dx &= \int_a^b (u'_{\min}(x))^2 + (u'_{\max}(x))^2 \, dx \\
&\leq \liminf_{i\to\infty} \int_a^b \left(u^i_{\min}{}'(x)\right)^2 + \left(u^i_{\max}{}'(x)\right)^2 \, dx \\
&= \liminf_{i\to\infty} \int_a^b |\mathrm{ap}Au^i(x)|^2 \, dx \\
&\leq 4 \liminf_{i\to\infty} \frac{\mathcal{G}^2(u^i(b), u^i(a))}{b-a} \\
&= 4 \liminf_{i\to\infty} \frac{|u^i_{\min}(b) - u^i_{\min}(a)|^2 + |u^i_{\max}(b) - u^i_{\max}(a)|^2}{b-a} \\
&= 4 \frac{|u_{\min}(b) - u_{\min}(a)|^2 + |u_{\max}(b) - u_{\max}(a)|^2}{b-a} \\
&= 4 \frac{\mathcal{G}^2(u(b), u(a))}{b-a}
\end{aligned}$$

where the first inequality follows from Lemma 23.7 in [17] and the second inequality is a consequence of Lemma 6. By Theorem 5, the multiple-valued function u is a Dirichlet 4-quasiminimizer. We denote the Cantor ternary set by

$$T = \cap_{i=1}^{\infty} T_i$$

where $T_1 = [0,1/3] \cup [2/3,1], T_2 = [0,1/9] \cup [2/9,1/3] \cup [2/3,7/9] \cup [8/9,1]$, etc. We can make the following observations:

(1) If $x \in T$ then $u^i_{\min}(x) = u^i_{\max}(x)$ for all $i \in \mathbb{N}_0$ hence $u_{\min}(x) = u_{\max}(x)$.
(2) If $x \notin T$ then there exists $i \in \mathbb{N}_0$ such that $x \notin T_j$ for each $j \geq i$. By construction, there exists a constant $\eta > 0$ such that $u^j_{\max}(x) - u^j_{\min}(x) = \eta$ for each $j \geq i$ hence $u_{\min}(x) \neq u_{\max}(x)$.

We deduce from (1) and (2) that

$$\{x \in [0,1] \mid \text{card}\,(\text{spt}(u(x))) = 1\} = T$$

hence the branch set of u is T since it contains no interval. □

It is clear that this construction can be adapted in order to obtain a quasiminimizer with a fat Cantor branch set.

2.3. **Application.** Let $a \in L^\infty(U^m(0,1), \mathbb{R})$ and consider the following functional

$$J(u; U^m(0,1)) = \int_{U^m(0,1)} a(x)\ |\text{ap}Au(x)|^2\ d\mathcal{L}^m x$$

defined for each $u \in \mathcal{Y}_2\left(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)\right)$. One says that $u : U^m(0,1) \to \mathbf{Q}_Q(\mathbb{R}^n)$ is $J-$minimizing if and only if $u \in \mathcal{Y}_2\left(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)\right)$ and, assuming u has boundary values $v \in \partial\mathcal{Y}_2\left(\partial U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)\right)$, one has

$$J\left(u; U^m(0,1)\right) = \inf\{J\left(w; U^m(0,1)\right) : w \in \mathcal{Y}_2\left(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)\right) \text{ has boundary values } v\}.$$

Theorem 8. *Let* $m \geq 2$ *be an integer. Assume* $v \in \partial\mathcal{Y}_2\left(\partial U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)\right)$ *and* $a(x) \geq \gamma > 0$ *for almost all* $x \in U^m(0,1)$. *Then there exists a strictly defined* $J-$*minimizing* $u \in \mathcal{Y}_2\left(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)\right)$ *that has boundary values* v *and*

(1) *if* $m = 2$, $u|_{B^2(0,\delta)}$ *is Hölder continuous for all* $0 < \delta < 1$;
(2) *if* $m \geq 3$ *and*

$$\frac{\|a\|_\infty}{\gamma} \leq \frac{1}{1 - \varepsilon_Q}$$

then $u|_{B^m(0,\delta)}$ *is Hölder continuous for all* $0 < \delta < 1$.

Proof. The existence proof is a direct adaptation of the proof of Theorem 2.2(2) in [2]. The regularity follows from the fact that u is a Dirichlet $\frac{\|a\|_\infty}{\gamma}$-quasiminimizer. Consequently it remains to apply Theorem 2 or Theorem 3. □

3. Dirichlet $\omega-$minimizer

Throughout this section $\omega : (0, \infty) \to \mathbb{R}$ will be a non-negative function.

Definition 9. *A strictly defined function* $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ *is a Dirichlet* ω*-minimizer if for every ball* $U^m(x,r) \subset U^m(0,1)$,

$$Dir(u; U^m(x,r)) \leq (1 + \omega(r))\ Dir(v; U^m(x,r))$$

where v *is a Dirichlet minimizing multiple-valued function having boundary values* $u|_{\partial U^m(x,r)} \in \partial\mathcal{Y}_2\left(\partial U^m(x,r), \mathbf{Q}_Q(\mathbb{R}^n)\right)$.

3.1. **Regularity.** The following regularity result is a direct consequence of Theorem 2 and Theorem 3.

Theorem 10. *Let* $m \geq 2$ *be an integer. Suppose that* $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ *is a strictly defined* ω*-minimizer such that* $Dir(u; U^m(0,1)) > 0$*. If* $\lim_{r\to 0} \omega(r) = 0$ *then*

$$u \in C^{0,\sigma}_{loc}(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$$

for some $0 < \sigma < 1$*.*

3.2. **Branch set of** $\omega-$**minimizer.** We will construct a one-dimensional Dirichlet $\omega-$minimizer with a unique branch point at the origin. First of all, we show that a one-dimensional single-valued $\omega-$minimizer does not have to be a straight line even if $\lim_{r\to 0} \omega(r) = 0$.

Proposition 11. $\sin x : (-\pi/4, \pi/4) \to \mathbb{R}$ *is a Dirichlet* $\omega-$*minimizer with*

$$\omega(r) = \frac{r^2}{\sin^2 r} - 1 \searrow 0 \text{ as } r \searrow 0.$$

Proof. It is easy to check that

$$\begin{aligned}
\mathrm{Dir}(\sin(\cdot); U^1(x,r)) &= \int_{x-r}^{x+r} (\cos s)^2 ds \\
&= \frac{\sin 2(x+r) - \sin 2(x-r)}{4} + r \\
&= \frac{\cos 2x \sin 2r}{2} + r
\end{aligned}$$

The Dirichlet minimizer v with boundary values $v(x-r) = \sin(x-r)$ and $v(x+r) = \sin(x+r)$ is the straight line with slope

$$\frac{\sin(x+r) - \sin(x-r)}{2r},$$

hence

$$\begin{aligned}
\mathrm{Dir}(v; U^1(x,r)) &= \left[\frac{\sin(x+r) - \sin(x-r)}{2r}\right]^2 2r \\
&= \frac{2\cos^2 x \sin^2 r}{r}.
\end{aligned}$$

Now it suffices to show that

$$\frac{\cos 2x \sin 2r}{2} + r \leq (1 + \omega(r)) \frac{2\cos^2 x \sin^2 r}{r}, \tag{4}$$

for any $U^1(x,r) \subset (-\pi/4, \pi/4)$. Define the function

$$W(x,r) = \left[\frac{\cos 2x \sin 2r}{2} + r\right] / \left[\frac{2\cos^2 x \sin^2 r}{r}\right].$$

Denote

$$W(x,r) = \frac{1}{4\sin^2 r} H(x,r),$$

where

$$H(x,r) = \frac{r\cos(2x)\sin(2r) + 2r^2}{\cos^2 x}.$$

We have

$$\frac{\partial H}{\partial x} = \frac{2r\sin x}{\cos^3 x}(2r - \sin(2r)).$$

For $r \in (0,1]$,

$$\frac{\partial H}{\partial x} \leq 0, \text{ for } x \in (-\pi/4, 0],$$

$$\frac{\partial H}{\partial x} \geq 0, \text{ for } x \in [0, \pi/4).$$

Since $W(x,r)$ is an even function of x, we conclude

$$W(x,r) \leq W(\pi/4, r) = \frac{r^2}{\sin^2 r}, \ \forall x \in (-\pi/4, \pi/4)$$

which proves (4). ☐

We are ready to construct a one-dimensional $\mathbf{Q}_2(\mathbb{R})-$valued $\omega-$minimizer with a branch point and $\lim_{r\to 0}\omega(r) = 0$.

Proposition 12. *The multiple-valued function*

$$u(x) = [[x]] + [[\sin x]] : (-\pi/4, \pi/4) \to \mathbf{Q}_2(\mathbb{R})$$

is a $\omega-$minimizer with

$$\omega(r) = \frac{r^2}{\sin^2 r} - 1 \searrow 0 \text{ as } r \searrow 0.$$

The origin is a branch point of u.

Proof. Proposition 12 is a direct consequence of Proposition 11 since the Dirichlet minimizer v with boundary values $u(x-r) = [[x-r]] + [[\sin(x-r)]], u(x+r) = [[x+r]] + [[\sin(x+r)]]$ is two straight lines connecting $x-r$ with $x+r$ and $\sin(x-r)$ with $\sin(x+r)$. ☐

Let us mention that the authors ignore the maximum Hausdorff dimension of branch sets of ω-minimizers.

4. DIRICHLET ALMOST MINIMIZER

Throughout this section $c \geq 0$ and $0 < \alpha < 1$ are real numbers.

Definition 13. *A strictly defined function $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ is a Dirichlet (c,α)-almost minimizer if for every ball $U^m(x,r) \subset U^m(0,1)$,*

$$Dir(u; U^m(x,r)) \leq Dir(v; U^m(x,r)) + cr^{m-2+\alpha}$$

where v is a Dirichlet minimizing multiple-valued function having boundary values $u|_{\partial U^m(x,r)} \in \partial\mathcal{Y}_2(\partial U^m(x,r), \mathbf{Q}_Q(\mathbb{R}^n))$.

4.1. **Regularity.** For the convenience of the reader, we shall recall a few results that we are going to need later.

Section 2.9 in [2]**.** *Corresponding to numbers $0 < s_0 < \infty$, $1 < K < \infty$, and (not necessarily distinct points) $q_1, \dots, q_Q \in \mathbb{R}^n$ we can find $J \in \{1, \dots, Q\}, k_1, \dots, k_J \in \{1, \dots, Q\}$, distinct points $p_1, \dots, p_J \in \{q_1, \dots, q_Q\}$, and $s_0 \le r \le Cs_0$ such that*

1. *$|p_i - p_j| > 2Kr$ for each $1 \le i < j \le J$,*
2. *$\mathcal{G}\left(\sum_{i=1}^{Q}[[q_i]], \sum_{i=1}^{J} k_i[[p_i]]\right) \le C(Q)s_0/(Q-1)^{1/2}$,*
3. *$z \in \mathbf{Q}_Q(\mathbb{R}^n)$ with $\mathcal{G}\left(z, \sum_{i=1}^{Q}[[q_i]]\right) \le s_0$ implies $\mathcal{G}\left(z, \sum_{i=1}^{J} k_i[[p_i]]\right) \le r$,*
4. *in case $J = 1$, diam$\Big($spt$\left(\sum_{i=1}^{Q}[[q_i]]\right)\Big) \le C(Q)s_0/(Q-1)$; here*

$$C(Q) = 1 + \left[(2K)(Q-1)^2\right]^1 + \left[(2K)(Q-1)^2\right]^2 + \dots + \left[(2K)(Q-1)^2\right]^{Q-1}.$$

Section 2.10 in [2]**.** *Corresponding to*

1. *$J \in \{1, 2, \dots, Q\}$,*
2. *$k_1, k_2, \dots, k_J \in \{1, 2, \dots, Q\}$ with $k_1 + k_2 + \dots + k_J = Q$,*
3. *distinct points $p_1, p_2, \dots, p_J \in \mathbb{R}^n$,*
4. *$0 < s_1 < s_2 = 2^{-1} \inf\{|p_i - p_j| : 1 \le i < j \le J\}$,*

we set

$$\mathbb{P} = \mathbf{Q}_Q(\mathbb{R}^n) \cap \{\sum_{i=1}^{Q}[[q_i]] : q_1, \dots, q_Q \in \mathbb{R}^n \text{ with } \\ \text{card}\{i : q_i \in B^n(p_j, s_1)\} = k_j \text{ for each } j = 1, \dots, J\}.$$

Then there exists a map $\Phi : \mathbf{Q}_Q(\mathbb{R}^n) \to \mathbb{P}$ such that

1. *$\Phi(q) = q$ whenever $q \in \mathbf{Q}_Q(\mathbb{R}^n)$ with $\mathcal{G}\left(q, \sum_{i=1}^{J} k_i[[p_i]]\right) \le s_1$,*
2. *$\Phi(q) = \sum_{j=1}^{J} k_j[[p_j]]$ whenever $q \in \mathbf{Q}_Q(\mathbb{R}^n)$ with $\mathcal{G}\left(q, \sum_{i=1}^{J} k_i[[p_i]]\right) \ge s_2$,*
3. *$\mathcal{G}(q, \Phi(q)) \le \mathcal{G}\left(q, \sum_{i=1}^{J} k_i[[p_i]]\right)$ for each $q \in \mathbf{Q}_Q(\mathbb{R}^n)$,*
4. *Lip $\Phi \le 1 + Q^{1/2}s_1/(s_2 - s_1)$.*

Theorem 2.12 in [2]**.** *For $z \in \mathbf{Q}_Q(\mathbb{R}^n)$ with $\mathcal{G}(z, q_0) > r$,*

1. *$\mathcal{G}(z, q) > s_0$.*
2. *$\mathcal{G}(z, q_0) \le \mathcal{G}(z, q) + \mathcal{G}(q, q_0) \le \mathcal{G}(z, q) + \dfrac{C(Q)s_0}{(Q-1)^{1/2}} \le [1 + C_2(Q)]\mathcal{G}(z, q)$, where $C_2(Q) = \dfrac{C(Q)}{(Q-1)^{1/2}}$.*
3. *$\mathcal{G}(z, q) \le \mathcal{G}(z, q_0) + \mathcal{G}(q_0, q) \le \mathcal{G}(z, q_0) + \dfrac{C(Q)s_0}{(Q-1)^{1/2}} \le [1 + C_2(Q)]\mathcal{G}(z, q_0)$.*

   Additionally, we define

   $$0 < s_0 \le r \le s_1 = K^{-1}s_2 < s_2 = 2^{-1}\inf\{|p_i - p_j| : 1 \le i < j \le J\}.$$

   Let $\Phi : \mathbf{Q}_Q(\mathbb{R}^n) \to \mathbf{Q}_Q(\mathbb{R}^n)$ be the semi-retraction mapping constructed in Section 2.10 in [2] *corresponding to $J, k_1, \dots, k_J, p_1, \dots, p_J, s_1, s_2$ above. For each $p \in \mathbf{Q}_Q(\mathbb{R}^n)$*

   $$\mathcal{G}(\Phi(p), p) \le \mathcal{G}(q_0, p) = 2\left[\mathcal{G}(q_0, p) - 2^{-1}\mathcal{G}(q_0, p)\right]$$

   so that

(4) $\mathcal{G}(\Phi(p), p) \leq 2[\mathcal{G}(p, q_0) - s_1/2]$.

Theorem A.1.6(17) in [2]. *Suppose* $f \in \mathcal{Y}_2(\partial U^m(0,1), \mathbb{R}^n), q \in V, 0 < r < \infty$, *and*

$$A = \partial B^m(0,1) \cap \{x : |f(x) - q| > r\}$$

with $\mathcal{H}^{m-1}(A) \leq m\alpha(m)/4$. *Then*

$$2^{-1-4/(m-1)}(2/\pi)^{2m-4+2/(m-1)}(m-1)\beta(m)^{-1} \int_A (|f(x) - q| - r)^2 \, d\mathcal{H}^{m-1} \leq \left[\mathcal{H}^{m-1}(A)\right]^{2/(m-1)} dir(f; A).$$

We are aiming to prove the following theorem:

Theorem 14. *Suppose* $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ *is a strictly defined Dirichlet* (c, α)*-almost minimizer such that* $Dir(u; U^m(0,1)) > 0$. *Then*

$$u \in C^{0,\sigma}_{loc}(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$$

for some $0 < \sigma < 1$.

We first prove an energy growth estimate for points with small normalized energy. This estimate is divided into two parts, on strong branch points (see definition below) and non-strong-branch points. Then we show that the energy density is zero for every point in the domain, which completes the interior regularity.

Definition 15. *For a multiple-valued function* u, *we define the strong branch set to be*

$$\begin{aligned} B_u = \{ & x \in U^m(0,1) : x \textit{ is a Lebesgue point of } \xi \circ u, \\ & \xi^{-1} \circ \rho \circ AV_{r,x}(\xi \circ u) = Q[[b_r]], \textit{ for any small enough radius } r > 0, b_r \in \mathbb{R}^n \}. \end{aligned}$$

where $AV_{r,x}(\xi \circ u) := \fint_{\partial U^m(x,r)} \xi \circ u$. *Obviously,* $x \in B_u$ *implies that* $u(x) = Q[[y]]$ *for some* $y \in \mathbb{R}^n$.

Lemma 16 (Hybrid Inequality). *There is a positive constant* C, *depending only on* m, n, Q, c, α *such that if* $0 < \lambda < 1$, $0 < \rho \leq 1$ *and* u *is a strictly defined* $(c, \alpha)-$*almost minimizer, then*

$$E_{\rho/2}(u) \leq \lambda E_\rho(u) + C \left[\rho^\alpha + \lambda^{-1} \rho^{-m} \int_{U^m(0,\rho)} |\xi \circ u - \mu|^2 dx \right], \tag{5}$$

for any constant vector $\mu \in \mathbb{R}^{PQ}$, *where* $E_r(u) = r^{2-m} Dir(u; U^m(0,r))$.

Proof. We first use Fubini's Theorem as in Section 2.3 in [11], to obtain a radius $\rho/2 \leq r \leq \rho$ so that $u|\partial U^m(0,r) \in \partial \mathcal{Y}_2(\partial U^m(0,r), \mathbf{Q}_Q(\mathbb{R}^n))$,

$$\int_{\partial U^m(0,r)} |\nabla_{\tan} u|^2 \, d\mathcal{H}^{m-1} \leq 8 \int_{U^m(0,\rho)} |Du|^2 dx, \tag{6}$$

$$\int_{\partial U^m(0,r)} |\xi \circ u - \mu|^2 d\mathcal{H}^{m-1} \leq 8 \int_{U^m(0,\rho)} |\xi \circ u - \mu|^2 dx. \tag{7}$$

Let $h : U^m(0,r) \to \mathbf{Q}_Q(\mathbb{R}^n)$ be Dirichlet minimizing with boundary values $u|\partial U^m(0,r)$.

$$\begin{aligned}\int_{U^m(0,r)} |Dh|^2 dx &= \int_{\partial U^m(0,r)} \left\langle \xi \circ h, \frac{\partial(\xi \circ h)}{\partial r} \right\rangle d\mathcal{H}^{m-1} \\ &= \int_{\partial U^m(0,r)} \left\langle \xi \circ h - \mu, \frac{\partial(\xi \circ h)}{\partial r} \right\rangle d\mathcal{H}^{m-1} \\ &\le \left[\int_{\partial U^m(0,r)} |\xi \circ h - \mu|^2 d\mathcal{H}^{m-1}\right]^{1/2} \left[\int_{\partial B^m(0,r)} \left|\frac{\partial h}{\partial r}\right|^2 d\mathcal{H}^{m-1}\right]^{1/2} \\ &= \left[\int_{\partial U^m(0,r)} |\xi \circ u - \mu|^2 d\mathcal{H}^{m-1}\right]^{1/2} \left[\int_{\partial U^m(0,r)} \left|\frac{\partial h}{\partial r}\right|^2 d\mathcal{H}^{m-1}\right]^{1/2}.\end{aligned}$$

By Section 2.6 in [2],

$$\int_{\partial U^m(0,r)} \left|\frac{\partial h}{\partial r}\right|^2 d\mathcal{H}^{m-1} \le \int_{\partial U^m(0,r)} |\nabla_{\tan} h|^2 d\mathcal{H}^{m-1} = \int_{\partial U^m(0,r)} |\nabla_{\tan} u|^2 d\mathcal{H}^{m-1}.$$

Therefore,

$$\int_{U^m(0,r)} |Dh|^2 dx \le \left[\int_{\partial U^m(0,r)} |\xi \circ u - \mu|^2 d\mathcal{H}^{m-1}\right]^{1/2} \left[\int_{\partial U^m(0,r)} |\nabla_{\tan} u|^2 d\mathcal{H}^{m-1}\right]^{1/2}$$

and

$$\begin{aligned}E_{\rho/2}(u) &= (\rho/2)^{2-m}\mathrm{Dir}(u; U^m(0,\rho/2)) \\ &\le (\rho/2)^{2-m}\mathrm{Dir}(u; U^m(0,r)) \\ &\le (\rho/2)^{2-m}\left[\int_{U^m(0,r)} |Dh|^2 dx + cr^{m-2+\alpha}\right] \\ &\le (\rho/2)^{2-m}\left[\int_{\partial U^m(0,r)} |\nabla_{\tan} u|^2 d\mathcal{H}^{m-1}\right]^{1/2} \left[\int_{\partial U^m(0,r)} |\xi \circ u - \mu|^2 d\mathcal{H}^{m-1}\right]^{1/2} \\ &\quad + 2^{m-2} c\rho^\alpha.\end{aligned}$$

Then we obtained the desired estimate by applying the inequality $ab \le \frac{1}{2}\delta a^2 + \frac{1}{2}\delta^{-1} b^2$, with $\delta = \frac{\lambda}{2^m}$ and using (6)(7) as follows

$$\begin{aligned}E_{\rho/2}(u) &\le \left(\frac{\rho}{2}\right)^{2-m}\left(\frac{1}{2}\delta \int_{\partial U^m(0,r)} |\nabla_{\tan} u|^2 d\mathcal{H}^{m-1} + \frac{1}{2}\delta^{-1}\int_{\partial U^m(0,r)} |\xi \circ u - \mu|^2 d\mathcal{H}^{m-1}\right) \\ &\quad + 2^{m-2} c\rho^\alpha \\ &\le \left(\frac{\rho}{2}\right)^{2-m}\left(4\delta \int_{U^m(0,\rho)} |Du|^2 dx + 4\delta^{-1}\int_{U^m(0,\rho)} |\xi \circ u - \mu|^2 dx\right) + 2^{m-2} c\rho^\alpha \\ &= \lambda E_\rho(u) + 4^m \lambda^{-1}\rho^{-m}\int_{U^m(0,\rho)} |\xi \circ u - \mu|^2 dx + 2^{m-2} c\rho^\alpha \\ &\le \lambda E_\rho(u) + C\left[\rho^\alpha + \lambda^{-1}\rho^{-m}\int_{U^m(0,\rho)} |\xi \circ u - \mu|^2 dx\right]\end{aligned}$$

where $C = \max\{2^{m-2}c, 4^m\}$. □

Let us introduce the family of functions that we will "blow up":

$$\mathcal{F} = \{u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n)) : u \text{ is } (c,\alpha) - \text{almost minimizing and } 0 \in B_u\}$$

Lemma 17 (Energy Improvement)**.** *There are positive constants ϵ_0, r_0, η and $\theta < 1$ so that, for any $(c,\alpha)-$almost minimizer $u \in \mathcal{F}$ with $E_{r_0}(u) < \epsilon_0^2$, one has*

$$E_{\theta r}(u) \leq \theta^{\omega_{2.13}} \max\{\eta r^\alpha, E_r(u)\}, \forall 0 < r < r_0, \tag{8}$$

where the constant $0 < \omega_{2.13} < 1$ is defined by Theorem 2.13 in [2].

Proof. If this theorem were false, then for any fixed positive $\theta < 1/2$, there would exist $(c,\alpha)-$almost minimizers $u_i \in \mathcal{F}$ and $r_i \to 0$ for which

$$\epsilon_i^2 := E_{r_i}(u_i) \to 0$$

$$r_i^{-\alpha} E_{\theta r_i}(u_i) \to \infty \tag{9}$$

as $i \to \infty$, but

$$E_{\theta r_i}(u_i) > \theta^{\omega_{2.13}} \epsilon_i^2 \tag{10}$$

for all i. The above relation (9) clearly implies that

$$\epsilon_i^{-2} r_i^\alpha \to 0, \text{ as } i \to \infty.$$

We define the blowing-up sequence $v_i : U^m(0,1) \to \mathbf{Q}_Q(\mathbb{R}^n)$ by

$$v_i(x) = \mu(\epsilon_i^{-1})_\sharp \circ \left[u_i(r_i x) - \xi^{-1} \circ \rho \circ AV_{r_i,0}(\xi \circ u_i)\right],$$

where the subtraction makes sense because $\xi^{-1} \circ \rho \circ AV_{r_i,0}(\xi \circ u_i) = Q[[b_i]]$ for some $b_i \in \mathbb{R}^n$.

The Dirichlet energy of v_i is clearly uniformly bounded by 1. As for the L^2 norms, we have

$$\begin{aligned}
&\int_{U^m(0,1)} \mathcal{G}^2\left(u_i(r_i x) - \xi^{-1} \circ \rho \circ AV_{r_i,0}(\xi \circ u_i), Q[[0]]\right) dx \\
&= \int_{U^m(0,1)} \mathcal{G}^2\left(u_i \circ \mu(r_i)(x) - \xi^{-1} \circ \rho \circ AV_{1,0}(\xi \circ u_i \circ \mu(r_i)), Q[[0]]\right) dx \\
&\leq \mathrm{Lip}(\xi^{-1})^2 \, \mathrm{Lip}(\rho)^2 \int_{U^m(0,1)} |\xi \circ u_i \circ \mu(r_i)(x) - AV_{1,0}(\xi \circ u_i \circ \mu(r_i))|^2 dx \\
&\leq C \, \mathrm{Lip}(\xi^{-1})^2 \, \mathrm{Lip}(\rho)^2 \, \mathrm{Dir}\left(\xi \circ u_i \circ \mu(r_i); U^m(0,1)\right) \\
&= C \, \mathrm{Lip}(\xi^{-1})^2 \, \mathrm{Lip}(\rho)^2 \, \epsilon_i^2,
\end{aligned}$$

where the second inequality comes from the Poincaré inequality (see Corollary 6.1 in [18]).

Using compactness theorem for multiple-valued functions (see Theorem 4.2 in [19]), there is a subsequence of v_i (still denoted as v_i) such that v_i converges weakly to $v \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$. Moreover, by similar argument as in Section 6.4 of [18], we can show that this convergence is actually strong and v is Dirichlet minimizing.

Let's estimate the extra term in the hybrid inequality

$$
\begin{aligned}
&\fint_{U^m(0,rr_i)} |\xi \circ u_i - AV_{r,0}(\xi \circ u_i \circ \mu(r_i))|^2 \, dx \\
&= \fint_{U^m(0,r)} |\xi \circ u_i(r_i x) - AV_{r,0}(\xi \circ u_i \circ \mu(r_i))|^2 \, dx \\
&\leq C r^{2-m} \int_{U^m(0,r)} |D(\xi \circ u_i \circ \mu(r_i))|^2 dx \quad \text{(by Poincaré inequality)} \\
&= C(\epsilon_i)^2 r^{2-m} \int_{U^m(0,r)} |Dv_i|^2 dx \\
&\leq C(\epsilon_i)^2 r^{2-m} \int_{U^m(0,r)} |Dv|^2 dx \quad \text{(by strong convergence)} \\
&\leq C(\epsilon_i)^2 r^{2-m} r^{m-2+2\omega_{2.13}} \mathrm{Dir}(v; U^m(0,1)) \quad \text{(interior regularity of Dirichlet minimizer)} \\
&= C(\epsilon_i)^2 r^{2\omega_{2.13}}.
\end{aligned}
$$

Applying the hybrid inequality to u_i with $\rho = 2\theta r_i$, we get

$$
\begin{aligned}
E_{\theta r_i}(u_i) &\leq \lambda E_{2\theta r_i}(u_i) + C\left[(2\theta r_i)^\alpha + \lambda^{-1} \fint_{U^m(0,2\theta r_i)} |\xi \circ u_i - AV_{2\theta,0}(\xi \circ u_i \circ \mu(r_i))|^2 \, dx\right] \\
&\leq \lambda E_{2\theta r_i}(u_i) + C\left[(2\theta r_i)^\alpha + \lambda^{-1} C(\epsilon_i)^2 (2\theta)^{2\omega_{2.13}}\right]
\end{aligned}
$$

Choosing a positive integer $k = k(\theta)$ for which $2^k\theta \leq 1 < 2^{k+1}\theta$, we iterate $k-1$ more times to obtain (we suppress all universal constant as C)

$$
\begin{aligned}
E_{\theta r_i}(u_i) &\leq \lambda^k E_{2^k \theta r_i}(u_i) + \sum_{j=1}^{k} \lambda^{j-1} C \left(2^j \theta r_i\right)^\alpha + \sum_{j=1}^{k} \lambda^{j-2} C(\epsilon_i)^2 \left(2^j \theta\right)^{2\omega_{2.13}} \\
&\leq \lambda^k 2^{m-2} \epsilon_i^2 + \sum_{j=1}^{\infty} \lambda^{j-1} C \left(2^j \theta r_i\right)^\alpha + \sum_{j=1}^{\infty} \lambda^{j-2} C \left(\epsilon_i\right)^2 \left(2^j \theta\right)^{2\omega_{2.13}} \\
&= \lambda^k 2^{m-2} \epsilon_i^2 + C(\theta r_i)^\alpha \frac{2^\alpha}{1-2^\alpha\lambda} + \frac{\lambda \cdot 2^{2\omega_{2.13}}}{1-\lambda\cdot 2^{2\omega_{2.13}}} C\lambda^{-2}\theta^{2\omega_{2.13}} (\epsilon_i)^2 \\
&= \left[\lambda^k 2^{m-2} + C\theta^\alpha \left(r_i^\alpha \epsilon_i^{-2}\right) \frac{2^\alpha}{1-2^\alpha\lambda} + \frac{\lambda \cdot 2^{2\omega_{2.13}}}{1-\lambda\cdot 2^{2\omega_{2.13}}} C\lambda^{-2}\theta^{2\omega_{2.13}}\right] (\epsilon_i)^2
\end{aligned}
$$

Taking $\lambda = \theta^{\frac{m+\omega_{2.13}}{k}}$, we have

$$
\lambda^k \cdot 2^{m-2} = \theta^{m+\omega_{2.13}} \cdot 2^{m-2} = \theta^m \cdot 2^{m-2} \cdot \theta^{\omega_{2.13}} \leq (1/2)^m \cdot 2^{m-2} \theta^{\omega_{2.13}} \leq \theta^{\omega_{2.13}}/4
$$

Since $\lambda = \theta^{\frac{m+\omega_{2.13}}{k}} \leq (2^{-k})^{\frac{m+\omega_{2.13}}{k}} = 2^{-(m+\omega_{2.13})}$,

$$
\begin{aligned}
\frac{\lambda \cdot 2^{2\omega_{2.13}}}{1-\lambda\cdot 2^{2\omega_{2.13}}} C\lambda^{-2}\theta^{2\omega_{2.13}} &\leq \frac{2^{2\omega_{2.13}} C}{1-2^{\omega_{2.13}-m}} \theta^{-\frac{m+\omega_{2.13}}{k}} \theta^{2\omega_{2.13}} \\
&\equiv M \theta^{\omega_{2.13} - \frac{m+\omega_{2.13}}{k}} \theta^{\omega_{2.13}},
\end{aligned}
$$

where $M = \dfrac{2^{2\omega_{2.13}} C}{1-2^{\omega_{2.13}-m}}$.

Let's choose θ small enough such that $\theta^{\omega_{2.13} - \frac{m+\omega_{2.13}}{k}} \leq 1/4M$. This is possible because it is equivalent to

$$\theta^{\omega_{2.13}} \le \theta^{\frac{m+\omega_{2.13}}{k}}/4M.$$

Notice that $\theta \ge 2^{-1-k}$, the right side of above one is greater than

$$2^{-(k+1)(m+\omega_{2.13})/k}/4M$$

which is bounded from below although when θ goes to zero, k goes to infinity. Notice that $\epsilon_i^{-2} r_i^{\alpha} \to 0$, as $i \to \infty$, for i sufficiently large enough,we have

$$E_{\theta r_i}(u_i) \le \left(\frac{1}{4}\theta^{\omega_{2.13}} + \frac{1}{4}\theta^{\omega_{2.13}} \right) \epsilon_i^2 < \theta^{\omega_{2.13}} \epsilon_i^2,$$

contradicting the choice of u_i in (10). $\square$

Iteration of (8) as Section 3.5 of [11] leads to the following energy decay estimate.

Theorem 18 (Energy decay for strong branch points). *If $u \in \mathcal{F}$ is (c,α)-almost minimizing, with $r_0^{2-m} \int_{U^m(0,r_0)} |Du|^2 \le \epsilon_0^2$, then*

$$r^{2-m} \int_{U^m(0,r)} |Du|^2 \le C r^{\alpha}, \ \textit{for } 0 \le r \le r_0$$

where ϵ_0 is as in the Energy Improvement.

Now we turn to non-strong-branch points and are going to prove an energy decay estimate by induction on Q. In particular, without loss of generality, we assume

$$\xi^{-1} \circ \rho \circ AV_{1,0}(\xi \circ u) \neq Q[[y]] \text{ for any } y \in \mathbb{R}^n.$$

Let $q^* = \rho \circ AV_{1,0}(\xi \circ u)$, $q = \xi^{-1}(q^*) = \sum_{i=1}^{Q} [[q_i]]$, and $q_0 = \sum_{i=1}^{J} k_i [[p_i]]$ is obtained from q using Section 2.9 in [2]. By the argument in Section 3 of [18], $J \ge 2$.

Lemma 19 (Construction of a comparison function). *Assume $m, n, Q \ge 2$ and $u \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ is a strictly defined $(c,\alpha)-$almost minimizer. Let $t_Q > 0$ be a real number such that*

$$\left[\frac{2Lip(\rho)Lip(\xi^{-1})\left(1 + C_2(Q)\right)}{s_0} \right]^2 Lip(\xi)^2 t_Q^{m-1} \le m\alpha(m)/4.$$

If u satisfies $dir(u; \partial U^m(0,1)) < t_Q^{m-1}$, then there is a comparison function $g \in \mathcal{Y}_2(U^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ satisfying the followings:

1. *$g = u$ on $\partial U^m(0,1)$,*
2. *$g|_{U^m(0,1-t_Q)} = \sum_{i=1}^{J} g_i$, where $g_i \in \mathcal{Y}_2(U^m(0,1-t_Q), \mathbf{Q}_{k_i}(\mathbb{R}^n))$ is Dirichlet minimizing and $\sum_{i=1}^{J} k_i = Q$,*
3. *$Dir(g; U^m(0,1) \sim U^m(0,1-t_Q)) \le \delta_Q \ dir(g; \partial U^m(0,1))$,*

for some constant δ_Q.

Proof. For each $x \in B^m(0,1) \sim B^m(0,1-t_Q)$, we define $\tau : B^m(0,1) \sim B^m(0,1-t_Q) \to \mathbb{R}$ and $F, G, H : B^m(0,1) \sim B^m(0,1-t_Q) \to \mathbb{R}^{PQ}$ by

$$\begin{aligned}
\tau(x) &= t_Q^{-1}(1 - |x|), \\
F(x) &= \xi \circ u(x/|x|), \\
H(x) &= \xi \circ \Phi \circ u(x/|x|), \\
G(x) &= (1 - \tau(x))F(x) + \tau(x)H(x).
\end{aligned}$$

Define $g : B^m(0,1) \sim B^m(0,1-t_Q) \to \mathbf{Q}_Q(\mathbb{R}^n)$ by

$$g|B^m(0,1) \sim B^m(0,1-t_Q) = \xi^{-1} \circ \rho \circ G|B^m(0,1) \sim B^m(0,1-t_Q).$$

On $\partial B^m(0,1)$, $g(x) = u(x)$. On $\partial B^m(0,1-t_Q)$, $g(x) = \Phi \circ u(x/|x|) = \sum_{i=1}^J h_i$, where $h_i \in \partial\mathcal{Y}_2(\partial B^m(0,1-t_Q), \mathbf{Q}_{k_i}(\mathbb{R}^n))$, for each $i = 1, \ldots, J$.
Let $g_i \in \mathcal{Y}_2(U^m(0,1-t_Q), \mathbf{Q}_{k_i}(\mathbb{R}^n))$ be a Dirichlet minimizing function with boundary h_i and $g|B^m(0,1-t_Q) = \sum_{i=1}^J g_i$. This completes (1) and (2).
By the definition of g,

$$\begin{aligned}
&\int_{U^m(0,1)\sim U^m(0,1-t_Q)} |\nabla g|^2 dx \\
&= \int_{U^m(0,1)\sim U^m(0,1-t_Q)} \left|\nabla(\xi^{-1} \circ \rho \circ G(x))\right|^2 dx \\
&\le \left[\mathrm{Lip}(\xi^{-1})\mathrm{Lip}(\rho)\right]^2 \int_{U^m(0,1)\sim U^m(0,1-t_Q)} |\nabla G(x)|^2 dx \\
&= \left[\mathrm{Lip}(\xi^{-1})\mathrm{Lip}(\rho)\right]^2 \left[\int_{U^m(0,1)\sim U^m(0,1-t_Q)} \left|\frac{\partial G}{\partial r}\right|^2 dx + \int_{U^m(0,1)\sim U^m(0,1-t_Q)} |\nabla_{\mathrm{Tan}} G(x)|^2 dx\right]
\end{aligned}$$

Also, by the definition of G, we can compute

$$\left|\frac{\partial G}{\partial r}\right| = \left|t_Q^{-1}(F(x) - H(x))\right| = \left|t_Q^{-1}\left(\xi \circ u(x/|x|) - \xi \circ \Phi \circ u(x/|x|)\right)\right|,$$

and

$$\begin{aligned}
\left|\nabla_{\partial U^m(0,|x|)} G(x)\right| &= |x|^{-1}\left|\nabla_{\partial U^m(0,1)} G(x)\right| \\
&= \left|\frac{1-\tau(x)}{|x|}\nabla_{\partial U^m(0,1)} F(x) + \frac{\tau(x)}{|x|}\nabla_{\partial U^m(0,1)} H(x)\right| \\
&\le \frac{1}{1-t_Q}\left|\nabla_{\partial U^m(0,1)}(\xi \circ u(x/|x|)) - \nabla_{\partial U^m(0,1)}(\xi \circ \Phi \circ u(x/|x|))\right| \\
&\le \frac{1}{1-t_Q}\mathrm{Lip}(\xi)\ \left(1 + \mathrm{Lip}(\Phi)\right)\left|\nabla_{\partial U^m(0,1)} u(x/|x|)\right|.
\end{aligned}$$

Therefore,

(11)

$$\begin{aligned}
&\int_{U^m(0,1)\sim U^m(0,1-t_Q)} |\nabla G(x)|^2 dx \\
&\le t_Q^{-2}\int_{U^m(0,1)\sim U^m(0,1-t_Q)} \left[\xi \circ u(x/|x|) - \xi \circ \Phi \circ u(x/|x|)\right]^2 dx \\
&+ (1-t_Q)^{-2}\left[\mathrm{Lip}(\xi)(1 + \mathrm{Lip}(\Phi))\right]^2 \int_{U^m(0,1)\sim U^m(0,1-t_Q)} \left|\nabla_{\partial U^m(0,1)}(u(x/|x|))\right|^2 dx \\
&\le t_Q^{-1}\int_{\partial U^m(0,1)} \left[\xi \circ u(x/|x|) - \xi \circ \Phi \circ u(x/|x|)\right]^2 d\mathcal{H}^{m-1} \\
&+ \frac{t_Q}{(1-t_Q)^2}\left[\mathrm{Lip}(\xi)(1 + \mathrm{Lip}(\Phi))\right]^2 \mathrm{dir}(u; \partial U^m(0,1)).
\end{aligned}$$

Now let us estimate

$$\int_{\partial U^m(0,1)} \left[\xi \circ u(x/|x|) - \xi \circ \Phi \circ u(x/|x|)\right]^2 d\mathcal{H}^{m-1}.$$

Let $Z = \partial U^m(0,1) \cap \{x : \mathcal{G}(u(x), q_0) > s_1\}$

$$\begin{aligned}
Z &\subset \partial U^m(0,1) \cap \left\{ x : \mathcal{G}(u(x), q) > \frac{s_1}{1 + C_2(Q)} \right\} \text{(by Theorem 2.12 in [2])} \\
&\subset \partial U^m(0,1) \cap \left\{ x : |\xi \circ u(x) - q^*| > \frac{s_1}{\mathrm{Lip}(\xi^{-1})(1 + C_2(Q))} \right\} \\
&\subset \partial U^m(0,1) \cap \left\{ x : |\xi \circ u(x) - q^*| > \frac{s_0}{2\mathrm{Lip}(\xi^{-1})(1 + C_2(Q))} \right\} \equiv A.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\mathcal{H}^{m-1}(A) \left(\frac{s_0}{2\mathrm{Lip}(\xi^{-1})(1 + C_2(Q))} \right)^2 \\
&\leq \int_{\partial U^m(0,1)} |\xi \circ u(x) - q^*|^2 d\mathcal{H}^{m-1} \\
&= \int_{\partial U^m(0,1)} |\xi \circ u(x) - \rho \circ AV_{1,0}(\xi \circ u)|^2 d\mathcal{H}^{m-1} \\
&\leq \mathrm{Lip}(\rho)^2 \int_{\partial U^m(0,1)} \left| \xi \circ u(x) - AV_{1,0}(\xi \circ u) \right|^2 d\mathcal{H}^{m-1}
\end{aligned} \tag{12}$$

and

$$\begin{aligned}
\mathcal{H}^{m-1}(Z) &\leq \mathcal{H}^{m-1}(A) \\
&\leq \left[\frac{2\mathrm{Lip}(\rho) \cdot \mathrm{Lip}(\xi^{-1})(1 + C_2(Q))}{s_0} \right]^2 \int_{\partial U^m(0,1)} \left| \xi \circ u(x) - AV_{1,0}(\xi \circ u) \right|^2 d\mathcal{H}^{m-1} \\
&\leq \left[\frac{2\mathrm{Lip}(\rho) \cdot \mathrm{Lip}(\xi^{-1})(1 + C_2(Q))}{s_0} \right]^2 \mathrm{Lip}(\xi)^2 \mathrm{dir}(u; \partial U^m(0,1)),
\end{aligned}$$

where the last inequality follows from [2] §A.1.6(3).
This permits us to estimate

$$\int_{\partial U^m(0,1)} |\xi\circ u - \xi\circ\Phi\circ u|^2 \, d\mathcal{H}^{m-1}$$

$$= \int_Z |\xi\circ u - \xi\circ\Phi\circ u|^2 \, d\mathcal{H}^{m-1}$$

$$\leq \mathrm{Lip}(\xi)^2 \int_Z [\mathcal{G}(u,\Phi\circ u)]^2 \, d\mathcal{H}^{m-1}$$

$$\leq 4\mathrm{Lip}(\xi)^2 \int_Z \left[\mathcal{G}(u(x),q_0) - \frac{s_1}{2}\right]^2 d\mathcal{H}^{m-1} (\text{by Theorem 2.12(4) in [2]})$$

$$\leq 4\mathrm{Lip}(\xi)^2 \int_Z \left[(1+C_2(Q))\ \mathcal{G}(u(x),q) - \frac{s_1}{2}\right]^2 d\mathcal{H}^{m-1}$$

(because on $Z, \mathcal{G}(u(x),q_0) > s_1 \geq s_0$ and Theorem 2.12(2) in [2])

$$\leq 4\left[\mathrm{Lip}(\xi)\mathrm{Lip}(\xi^{-1})\,(1+C_2(Q))\right]^2 \int_Z \left[|\xi\circ u(x) - q^*| - \frac{s_1}{2\mathrm{Lip}(\xi^{-1})\,(1+C_2(Q))}\right]^2 d\mathcal{H}^{m-1}$$

$$\leq 4\left[\mathrm{Lip}(\xi)\mathrm{Lip}(\xi^{-1})(1+C_2(Q))\right]^2 \int_Z \left[|\xi\circ u(x) - q^*| - \frac{s_0}{2\mathrm{Lip}(\xi^{-1})\,(1+C_2(Q))}\right]^2 d\mathcal{H}^{m-1}$$

(because $s_0 \leq s_1$)

$$\leq 4\left[\mathrm{Lip}(\xi)\mathrm{Lip}(\xi^{-1})(1+C_2(Q))\right]^2 C_1 \left[\mathcal{H}^{m-1}(A)\right]^{2/(m-1)} \mathrm{dir}(\xi\circ u; A)$$

(by Theorem A.1.6(17) in [2] and the estimate of $\mathcal{H}^{m-1}(A)$ in (12))

$$\left(C_1 = [2^{-1-4/(m-1)}(2/\pi)^{2m-4+2/(m-1)}(m-1)\beta(m)^{-1}]^{-1}\right)$$

$$\leq C\,(\xi,\Phi,m,Q)\,\mathrm{dir}\,(u;\partial U^m(0,1))^{1+\dfrac{2}{m-1}}$$

In summary, we have

$$\begin{aligned}
&\int_{U^m(0,1)\sim U^m(0,1-t_Q)} |\nabla g|^2 \, dx \\
&\leq C(\xi,\Phi,m,n,Q)\left[t_Q^{-1}\mathrm{dir}(g;\partial U^m(0,1))^{1+\dfrac{2}{m-1}} + t_Q\mathrm{dir}(g;\partial U^m(0,1))\right] \\
&= C(\xi,\Phi,m,n,Q)\left[t_Q^{-1}\mathrm{dir}(g;\partial U^m(0,1))^{\dfrac{2}{m-1}} + t_Q\right]\mathrm{dir}(g;\partial U^m(0,1)) \\
&\leq 2C(\xi,\Phi,m,n,Q)\ t_Q\ \mathrm{dir}(g;\partial U^m(0,1)) \\
&= \delta_Q\ \mathrm{dir}(g;\partial U^m(0,1)),
\end{aligned}$$

where $\delta_Q = 2C(\xi,\Phi,m,n,Q)t_Q$. □

Remark. (1) *Here is the scaled version: Assume* $(c,\alpha)-$*almost minimizer* $u \in \mathcal{Y}_2(U^m(0,r),\mathbf{Q}_Q(\mathbb{R}^n))$ *satisfies*

$$dir(u;\partial U^m(0,r)) < r^{m-3}t_Q^{m-1},$$

then the constructed comparison function g *with* $g = u$ *on* $\partial U^m(0,r)$ *satisfies*

$$Dir(g;U^m(0,r)\sim U^m(0,r(1-t_Q))) \leq \delta_Q r\, dir(u;\partial U^m(0,r)).$$

(2) *The smallness assumption of* $dir(u;\partial U^m(0,r))$ *can be replaced by the smallness of* $Dir(u;U^m(0,r))$. *This is because we choose* $g\in\mathcal{Y}_2(U^m(0,r),\mathbf{Q}_Q(\mathbb{R}^n))$ *with* $g=u$ *on* $\partial U^m(0,r)$. *By the squeeze formula ([2] §2.6), we have*

$$\begin{aligned}2r\,dir(u;\partial U^m(0,r)) &= 2r\,dir(g;\partial U^m(0,r))\\ &=(m-2)Dir(g;U^m(0,r))+r\,Dir(g;U^m(0,r))'\\ &\le (m-2)Dir(u;U^m(0,r))+r\,Dir(g;U^m(0,r))'\end{aligned}$$

So in the blowing-up analysis, by choosing a good slicing by $\partial U^m(0,r)$ *and rescaling, we can get small* $Dir(g;U^m(0,r))'$ *and hence the smallness of* $dir(u;\partial U^m(0,r))$.

Theorem 20 (Energy decay for non-strong-branch points). *Let* $m,n\ge 2$. *There exists a small number* $\epsilon=\epsilon(Q,m,n,c,\alpha)$ *such that any* $(c,\alpha)-$*almost minimizer* $u\in\mathcal{Y}_2(U^m(0,1),\mathbf{Q}_Q(\mathbb{R}^n))$ *with* $0\notin B_v$ *such that* $r_0^{2-m}Dir(u;U^m(0,r_0))<\epsilon$ *for some* $0<r_0\le 1$ *satisfies*

$$Dir(u;U^m(0,r))\le Cr^{m-2+\beta},0<r\le r_0$$

for some universal positive constants C,β.

Proof. Let g be the comparison function on $B^m(0,r)$,

$$\begin{aligned}\mathrm{Dir}(u;U^m(0,r)) &\le \mathrm{Dir}(g,U^m(0,r(1-t_Q)))+\delta_Q r\mathrm{dir}(g;\partial U^m(0,r))+cr^{m-2+\alpha}\\ &=\mathrm{Dir}(g;U^m(0,r(1-t_Q)))+\delta_Q r\mathrm{dir}(u;\partial U^m(0,r))+cr^{m-2+\alpha}\\ &\le \mathrm{Dir}(g,U^m(0,r(1-t_Q)))+\delta_Q r\mathrm{Dir}(u,U^m(0,r))'+cr^{m-2+\alpha}\end{aligned}$$

Applying the interior regularity of Dirichlet minimizing function to $g|B^m(0,r(1-t_Q))$,

$$\mathrm{Dir}(u;U^m(0,r))\le r^{m-2+2\omega_{2.13}}+\delta_Q r\mathrm{Dir}(u;U^m(0,r))'+cr^{m-2+\alpha}.$$

Denote $\omega=\min\{2\omega_{2.13},\alpha\}, N=1+c, D(r)=\mathrm{Dir}(u;U^m(0,r))$, we have

$$D(r)\le\delta_Q rD'(r)+Nr^{m-2+\omega},$$

i.e.

$$D'-\frac{D}{\delta_Q r}+\frac{N}{\delta_Q}r^{m-3+\omega}\ge 0.$$

Multiplying $r^{\frac{-1}{\delta_Q}}$,

$$\frac{d}{dr}\left[D(r)r^{\frac{-1}{\delta_Q}}+\frac{N}{\delta_Q\left(m-2+\omega-\frac{1}{\delta_Q}\right)}r^{m-2+\omega-\frac{1}{\delta_Q}}\right]\ge 0.$$

Therefore,

$$D(r)r^{\frac{-1}{\delta_Q}}+\frac{N}{\delta_Q\left(m-2+\omega-\frac{1}{\delta_Q}\right)}r^{m-2+\omega-\frac{1}{\delta_Q}}\le M:=D(1)+\frac{N}{\delta_Q\left(m-2+\omega-\frac{1}{\delta_Q}\right)}$$

$$D(r)\le Mr^{\frac{1}{\delta_Q}}+\frac{N}{1-\delta_Q(m-2+\omega)}r^{m-2+\omega}$$

Letting $C = \max\left\{M, \frac{N}{1-\delta_Q(m-2+\omega)}\right\}, \beta = \min\left\{\frac{1}{\delta_Q} - (m-2), \omega\right\}$ (which is positive since we can choose δ_Q small enough) finishes the proof. □

In spirit of Morrey's growth lemma, it only remains to show Theorem 21 to prove Theorem 14.

Theorem 21. *For any $(c, \alpha)-$almost minimizer u,*

$$\liminf_{r \downarrow 0} r^{2-m} Dir(u; U^m(x,r)) = 0, \forall x \in U^m(0,1).$$

Remark. *It is well-known (e.g* [13] *lemma 2.1.1) that the above one holds $\mathcal{H}^{m-2}$ a.e. Also, it suffices to prove this for $x = 0$.*

Lemma 22 (Monotonicity Formula)**.** *If u is a $(c,\alpha)-$almost minimizer, then for any $0 < t < s < 1$, we have*

$$\int_t^s r^{2-m} \int_{\partial U^m(0,r)} \left|\frac{\partial u}{\partial r}\right|^2 d\mathcal{H}^{m-1} dr - \frac{(m-2)c}{\alpha}(s^\alpha - t^\alpha)$$
$$\leq s^{2-m} Dir(u; U^m(0,s)) - t^{2-m} Dir(u; U^m(0,t)).$$

In particular, $r^{2-m} Dir(u; U^m(0,r)) + \frac{(m-2)c}{\alpha} r^\alpha$ is nondecreasing in r. Hence the limiting density

$$\Theta_u(0) := \lim_{r\downarrow 0} r^{2-m} \int_{U^m(0,r)} |Du|^2 dx$$

exists.

Proof. For any $0 < r < 1$, define a function $g : B^m(0,1) \to \mathbf{Q}_Q(\mathbb{R}^n)$ by setting

$$g(x) = \begin{cases} f\left(r\frac{x}{|x|}\right) & |x| \leq r \\ f(x) & \text{otherwise.} \end{cases}$$

By definition of almost minimizer, we have

$$\begin{aligned} &\int_{U^m(0,r)} |Du|^2 dx \\ &\leq cr^{m-2+\alpha} + \int_{U^m(0,r)} |Dg|^2 dx \\ &= cr^{m-2+\alpha} + \frac{r}{m-2}\int_{\partial U^m(0,r)} |\nabla_{\tan} u|^2 \, d\mathcal{H}^{m-1} \\ &= cr^{m-2+\alpha} + \frac{r}{m-2}\left[\frac{d}{d\rho}|_{\rho=r} \int_{U^m(0,\rho)} |Du|^2 dx - \int_{\partial U^m(0,r)} \left|\frac{\partial u}{\partial r}\right|^2 d\mathcal{H}^{m-1}\right] \end{aligned}$$

Multiplying by $(m-2)r^{1-m}$ to both sides,

$$r^{2-m}\int_{\partial U^m(0,r)} \left|\frac{\partial u}{\partial r}\right|^2 d\mathcal{H}^{m-1} - (m-2)cr^{\alpha-1} \leq \frac{d}{d\rho}|_{\rho=r}(\rho^{2-m}\mathrm{Dir}(u; U^m(0,\rho))).$$

Then integrate this from t to s. □

From the monotonicity formula, following the standard technique (e.g. [16]), we obtain a tangent map $v \in \mathcal{Y}_2(B^m(0,1), \mathbf{Q}_Q(\mathbb{R}^n))$ of u at the origin. v is a homogeneous degree zero Dirichlet minimizing function with

$$\theta_v(0) = \theta_u(0).$$

From the interior regularity, v has to be constant hence $\theta_u(0) = \theta_v(0) = 0$. This finishes the proof of Theorem 21 and also the proof of Theorem 14.

4.2. **Branch set of almost minimizer.** We will now produce a Dirichlet almost minimizer with a fractal branch set. We begin with some definitions:

1. A *losange* above an interval $I = [a,b] \subset \mathbb{R}$ is a multiple-valued function $u : [a,b] \to \mathbf{Q}_2(\mathbb{R})$ whose the graph is a parallelogram with the following vertices $[a,0], [b,0], [(a+b)/2, (b-a)/2], [(a+b)/2, (a-b)/2]$.
2. A *pluri-losange* is a multiple-valued function $u : [0,1] \to \mathbf{Q}_2(\mathbb{R})$ which admits a partition of $[0,1]$ in intervals $\{I_j\}$ such that u is a losange above I_j or a map of the form $x \to 2[[0]]$ above I_j for each j.

Figure 2 gives examples of pluri-losanges. It is clear that a pluri-losange u is Lipschitz with $\mathrm{Lip}(u) \leq \sqrt{2}$ and one readily checks that a pluri-losange is a Dirichlet $(2, 1/2)-$almost minimizer. We also notice that a pluri-losange is not a Dirichlet quasiminimizer or an ω-minimizer.

We will now create a particular sequence of pluri-losanges $\{u^i\}$. We define $u^1 : [0,1] \to \mathbf{Q}_2(\mathbb{R})$ by requiring that

- u^1 is a pluri-losange,
- $u^1(1/3) = 0$,
- u^1 is a losange only above $[1/3, 2/3]$.

We define $u^2 : [0,1] \to \mathbf{Q}_2(\mathbb{R})$ by requiring that

- u^2 is a pluri-losange,
- $u^2(1/3) = 0$,
- u^2 is a losange only above the intervals $[1/3, 2/3], [1/9, 2/9]$ and $[7/9, 8/9]$.



FIGURE 2. The pluri-losanges u^1, u^2 and u^3.

The reader will easily imagine how the remaining terms of the sequence are defined. By mimicking the arguments used in Section 2.2, we can extract a subsequence which converges uniformly to a Lipschitz $(2, 1/2)-$almost minimizer $u : [0,1] \to \mathbf{Q}_2(\mathbb{R})$ whose the branch set is the Cantor ternary set. This construction can also be adapted in order to obtain an almost minimizer with a fat Cantor branch set.

5. Acknowledgements

The authors are thankful to Thierry De Pauw and Robert Hardt for suggesting the problem and helpful discussions.

References

[1] F. J. Almgren, Jr., *Dirichlet's problem for multiple valued functions and the regularity of mass minimizing integral currents*, Minimal Submanifolds and Geodesics (M. Obata, editor), Kaigai, Tokyo, 1978, 1-6.

[2] F. J. Almgren, Jr., *Almgren's big regularity paper.* **Q***-valued functions minimizing Dirichlet's integral and the regularity of area-minimizing rectifiable currents up to codimension 2.* With a preface by Jean E. Taylor and Vladimir Scheffer. World Scientific Monograph Series in Mathematics, 1. World Scientific Publishing Co., Inc., River Edge, NJ, 2000.

[3] G. Anzellotti, *On the* $C^{1,\alpha}$ *regularity of* ω*-minima of quadratic functionals*, Boll. U.M.I., (VI) 2 (1983), 195-212.

[4] L. C. Evans R. F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992.

[5] H. Federer, *Geometric measure theory*, Springer-Verlag, Berlin and New York, 1969.

[6] M. Giaquinta, *Multiple integrals in calculus of variations and nonlinear elliptic systems*, Annals of Math. Studies, 105 Princeton Univ. Press, 1983.

[7] M. Giaquinta, E. Giusti, *Quasi Minima*, Ann. Inst. H. Poincaré (Analyse non lineaire), 1 (1984), 79-107.

[8] E. Giusti, *Metodi diretti nel calcolo delle variazioni*, U.M.I. 1994.

[9] J. Goblet, *A selection theory for multiple-valued functions in the sense of Almgren*, Ann. Aca. Scient. Fenn. Mat., Vol. 31, 2006, 297-314.

[10] R. Hardt, D. Kinderlehrer, *Some regularity results in ferromagnetism*, Commun. in Partial Differential Equations, 25 (7&8), 1235-1258 (2000)

[11] R. Hardt, D. Kinderlehrer, F.H. Lin, *Existence and partial regularity of static liquid crystal configurations*, Commun. Math. Phys. 105, 547-570 (1986)

[12] C.C. Lin, *Variational problems with multiple-valued functions and mappings*, PhD Thesis, Rice University

[13] F.H. Lin, X.Y. Yang, *Geometric measure theory: an Introduction*, International Press, 2003

[14] P. Mattila, *Lower semicontinuity, existence and regularity theorems for elliptic variational integrals of multiple valued functions*, Trans. Amer. Math. Soc. 280 (1983), no. 2, 589-610.

[15] C. B. Morrey, Jr., *Multiple integrals in the calculus of Variations*, Springer-Verlag, Berlin and New york, 1966.

[16] L. Simon, *Theorems on regularity and singularity of energy minimizing maps*, Birkhäuser, 1996

[17] M. Willem, *Analyse fonctionnelle élémentaire*, Cassini, Paris, 2003.

[18] W. Zhu, *A regularity theory for multiple-valued Dirichlet minimizing maps*, math.OC/0608178

[19] W. Zhu, *An energy reducing flow for multiple-valued functions*, arXiv:math.AP/0606478

Jordan Goblet, Département de mathématique, Université catholique de Louvain, Chemin du cyclotron 2, 1348 Louvain-La-Neuve (Belgium)

E-mail address: `goblet@math.ucl.ac.be`

Wei Zhu, Department of Mathematics, Rice University, 6100 Main Street, Houston, TX 77005 (U.S.A)

E-mail address: `weizhu@rice.edu`